SECTION H DRUG/ALCOHOL USE - ASI

*H8. Cocaine

(regular use can = 2 day binges/week)

Directions:	How many days in the past 30 before detox., have you used? How many years in your life have you regularly used? (3x/week or more)					
	Route of Administration Types:					
		0. Not applicable				
	1. Oral					
	2. Nasal					
	3. Smoking 4. Non-IV Injection					
	5. IV					
Note the usual o	or most recent route. For n	nore than one route, choose the	most severe. The route	es are listed from		
least severe to m		ask 1st	≥3x/week			
		(# of days)	(# of years)	Route of		
		Past 30 Days	Lifetime	Admin		
		1				
*H1. Alcohol	(any use at all)					
	1					
*H2. Alcohol	(to intoxication)					
(3 or me	ore &/or 'feel the effe	ects')				
(regular	use can = 2 day binges	s/week)				
*H3. Heroin						
*H4. Methado	ne					
*H5. Other Op	piates/Analgesics					
*H6. Barbitura	ntes					
H7 Cadative	s/Hypnotics/Tranquiliz	OTC .				

How many years in	your life have you regula	rly used? (3x/we	ek or more
How do you use it?			
Route of Administration	Types:		
0. Not applicable			
1. Oral			
2. Nasal			
3. Smoking			
4. Non-IV Injection			
5. IV			
Note the usual or most recent route. For n	nore than one route, choose th	e most severe. The route	es are listed fro
least severe to most severe.			o an o rainca yr c
	ask 1st	≥3x/week	
	(# of days) Past 30 Days	(# of years) Lifetime	Route of
	<i>[</i>		
*H9. Amphetamines	7 3 5 mile		
* 1			
*H10. Marijuana/Cannabis			-
*H11. Hallucinogens			
*H12. Inhalants			<u> </u>

How many days in the past 30 before detox., have you used....?

Directions:

*H13. More than 1 substance per day (may include alcohol)

(Interviewer should determine the major drug of number next to the drug in questions 01-12, "00	abuse. Code the " = no problem,	(00-16)
"15" = alcohol & one or more drugs, "16" = more Ask patient when not clear.)	re than one drug.	
W. C. W		
H15. How many times have you had:	THE ALLEDER	
	H15a. Alcohol DT's?	time
	H15b. Overdosed on Drugs?	time
(Delirium Tremens (DT's): occur 24-48 hours aft alcohol intake, shaking, severe disorientation, feve medical attention.)		
(Overdoses (OD): Requires intervention by some Include suicide attempts by OD.)	cone to recover, not simply sleeping it off.	
*H16. How much money would you say you sp	ent during the past 30 days (before d	letox.) on:
	H16a. Alcohol? \$	
	H16b. Drugs? \$	
(Only count actual money spent. What is the find		
*H17. How many days in the past 30 (before de	tov) have you experienced.	
1117. How many days in the past 30 (before de	H17a. Alcohol Problems?	(0-30)
	H17b. Drug Problems?	(0-30)
(Include only: craving, withdrawal symptoms, dis		(0-30)
to stop and being unable to.)		
For questions H18 and H19, ask the patient to use the Pa additional substance abuse treatment. (Show Card 16)	tient Rating Scale. The patient is rating the	need for
0. Not at All		
1. Slightly		
Moderately Considerably		
4. Extremely	2	
*TITO IT tblad on both and bosses sons boom	in the next 20 days (hefers datas) by	.4
*H18. How troubled or bothered have you been		
	H18a. Alcohol Problems?	
	H18b. Drug Problems?	(0-4)
*H19. How important to you now is treatment for	or these:	
	H19a. Alcohol Problems?	(0-4)
	H19b. Drug Problems?	

.

E ... (12)

: E