

# SDM4 in R: Regression Wisdom (Chapter 8)

Nicholas Horton ([nhorton@amherst.edu](mailto:nhorton@amherst.edu))

January 2, 2017

## Introduction and background

This document is intended to help describe how to undertake analyses introduced as examples in the Fourth Edition of *Stats: Data and Models* (2014) by De Veaux, Velleman, and Bock. More information about the book can be found at [http://wps.aw.com/aw\\_deveaux\\_stats\\_series](http://wps.aw.com/aw_deveaux_stats_series). This file as well as the associated R Markdown reproducible analysis source file used to create it can be found at <http://nhorton.people.amherst.edu/sdm4>.

This work leverages initiatives undertaken by Project MOSAIC (<http://www.mosaic-web.org>), an NSF-funded effort to improve the teaching of statistics, calculus, science and computing in the undergraduate curriculum. In particular, we utilize the `mosaic` package, which was written to simplify the use of R for introductory statistics courses. A short summary of the R needed to teach introductory statistics can be found in the `mosaic` package vignettes (<http://cran.r-project.org/web/packages/mosaic>).

## Chapter 8: Regression Wisdom

### Section 8.1: Examining residuals

Figure 8.1 (page 220) displays the scatterplot of heart rate vs duration for the Penguins dataset (along with a superimposed regression line and a smoother).

```
library(mosaic); library(readr)
options(digits=3)
Penguins <- read_csv("http://nhorton.people.amherst.edu/sdm4/data/Penguins.csv")
xyplot(DiveHeartRate ~ Duration, ylab="Dive Heart Rate (bpm)",
       xlab="Duration (mins)", type=c("p", "r", "smooth"), data=Penguins)
```




Figure 8.2 (page 220) displays the residuals from a linear regression model as a function of duration.

```
Penguinmod <- lm(DiveHeartRate ~ Duration, data=Penguins)
xyplot(resid(Penguinmod) ~ Duration, type=c("p", "r", "smooth"), data=Penguins)
```




Figure 8.3 (page 221) displays the histogram of residuals for the cereal data from Chapter 7.

```
Cereals <- read_csv("http://nhorton.people.amherst.edu/sdm4/data/Cereals.csv")
Cerealmod <- lm(calories ~ sugars, data=Cereals)
histogram(~ resid(Cerealmod), width=7.5, center=3.75)
```





Figure 8.4 (page 221) displays a scatterplot of the residuals vs predicted values. Without jittering, the display has an odd pattern.

```
xyplot(resid(Cerealmod) ~ fitted(Cerealmod))
```



By adding some random noise we can more easily observe values that are shared by more than one cereal.

```
xyplot(jitter(resid(Cerealmod)) ~ jitter(fitted(Cerealmod)))
```




Figure 8.5 (page 222) displays the scatterplot stratified by what shelf it is displayed on at the store.


```
tally(~ shelf, data=Cereals)
```

```
## shelf
## 1 2 3
## 20 21 36

Cereals <- mutate(Cereals, shelfgrp = derivedFactor(
  bottomshelf = shelf==1,
  middleshelf = shelf==2,
  topshelf = shelf==3
))
tally(~ shelfgrp, data=Cereals)

## shelfgrp
## bottomshelf middleshelf      topshelf
##          20            21            36
```

```
xyplot(calories ~ sugars, group=shelfgrp, auto.key=TRUE,  
lwd=2, type=c("p", "r"), data=Cereals)
```



Section 8.2: Extrapolation and reaching beyond the data

Section 8.3: Outliers, leverage, and influence

Section 8.4: Lurking variables and causation

Section 8.5: Working with summary values