

SDM4 in R: Testing Hypotheses about Proportions (Chapter 19)

Nicholas Horton (nhorton@amherst.edu)

January 2, 2017

Introduction and background

This document is intended to help describe how to undertake analyses introduced as examples in the Fourth Edition of *Stats: Data and Models* (2014) by De Veaux, Velleman, and Bock. More information about the book can be found at http://wps.aw.com/aw_deveaux_stats_series. This file as well as the associated R Markdown reproducible analysis source file used to create it can be found at <http://nhorton.people.amherst.edu/sdm4>.

This work leverages initiatives undertaken by Project MOSAIC (<http://www.mosaic-web.org>), an NSF-funded effort to improve the teaching of statistics, calculus, science and computing in the undergraduate curriculum. In particular, we utilize the `mosaic` package, which was written to simplify the use of R for introductory statistics courses. A short summary of the R needed to teach introductory statistics can be found in the `mosaic` package vignettes (<http://cran.r-project.org/web/packages/mosaic>).

Chapter 19: Testing hypotheses for proportions

Section 19.1: Hypotheses


We can reproduce the calculation in Figure 19.1 (page 495).

```
sdp <- sqrt(.2*.8/400); sdp
```

```
## [1] 0.02
```

```
xpnorm(0.17, mean=0.20, sd=sdp)
```

```
##  
## If  $X \sim N(0.2, 0.02)$ , then  
##  
##  $P(X \leq 0.17) = P(Z \leq -1.5) = 0.0668$   
##  $P(X > 0.17) = P(Z > -1.5) = 0.9332$ 
```



```
## [1] 0.0668
```

```
zval <- (0.17 - 0.20)/sdp; zval
```

```
## [1] -1.5
```

```
pnorm(zval, mean=0, sd=1)
```

```
## [1] 0.0668
```

Section 19.3: Reasoning of hypothesis testing

The “For Example (page 499)” lays out how to find a p-value for the one proportion z-test.

```
y <- 61; n <- 90; phat <- y/n; phat
```


```
## [1] 0.678
```

```
nullp <- 0.8  
sdp <- sqrt(nullp*(1-nullp)/n); sdp
```

```
## [1] 0.0422
```

```
onesidep <- xpnorm(phat, mean=nullp, sd=sdp); onesidep
```

```
##  
## If  $X \sim N(0.8, 0.0422)$ , then  
##  
##  $P(X \leq 0.678) = P(Z \leq -2.9) = 0.00187$   
##  $P(X > 0.678) = P(Z > -2.9) = 0.9981$ 
```



```
## [1] 0.00187
```

```
twosidep <- 2*onesidep; twosidep
```

```
## [1] 0.00375
```

or we can carry out the exact test (not described by the book):

```
binom.test(y, n, p=nullp)
```

```
##  
##  
##  
## data: y out of n  
## number of successes = 60, number of trials = 90, p-value = 0.006  
## alternative hypothesis: true probability of success is not equal to 0.8  
## 95 percent confidence interval:  
## 0.571 0.772  
## sample estimates:  
## probability of success  
## 0.678
```