

SDM4 in R: Randomness and Probability (Chapter 15)

Nicholas Horton (nhorton@amherst.edu)

August 12, 2017

Introduction and background

This document is intended to help describe how to undertake analyses introduced as examples in the Fourth Edition of *Stats: Data and Models* (2014) by De Veaux, Velleman, and Bock. More information about the book can be found at http://wps.aw.com/aw_deveaux_stats_series. This file as well as the associated R Markdown reproducible analysis source file used to create it can be found at <http://nhorton.people.amherst.edu/sdm4>.

This work leverages initiatives undertaken by Project MOSAIC (<http://www.mosaic-web.org>), an NSF-funded effort to improve the teaching of statistics, calculus, science and computing in the undergraduate curriculum. In particular, we utilize the `mosaic` package, which was written to simplify the use of R for introductory statistics courses. A short summary of the R needed to teach introductory statistics can be found in the `mosaic` package vignettes (<http://cran.r-project.org/web/packages/mosaic>). A paper describing the `mosaic` approach was published in the *R Journal*: <https://journal.r-project.org/archive/2017/RJ-2017-024>.

Chapter 15: Randomness and Probability

Section 15.1: Center (the Expected Value)

We can replicate the calculation on page 390:

```
library(mosaic); library(readr); options(digits=3)
x <- c(10000, 5000, 0)
prob <- c(1/1000, 2/1000, 997/1000)
sum(prob)      # sums to 1
```

```
## [1] 1
```

```
expect <- sum(x*prob); expect # expected value
```

```
## [1] 20
```

Section 15.2: Spread (The Standard Deviation)

We can continue with the example from page 392:

```
xminmu <- x - expect; xminmu
```

```
## [1] 9980 4980 -20
```

```
myvar <- sum(xminmu^2*prob); myvar
```

```
## [1] 149600
```

```
sd <- sqrt(myvar); sd
```

```
## [1] 387
```

Section 15.3: Shifting and Combining Random Variables

Let's replicate the values from the example on page 394:

```
ex <- 5.83; varx <- 8.62^2  
ed <- ex+5; ed
```

```
## [1] 10.8
```

```
vard <- varx; vard
```

```
## [1] 74.3
```

```
sqrt(vard)
```

```
## [1] 8.62
```

Section 15.5: Continuous random variables

Let's replicate Figure 15.1 (page 400):

```
xpnorm(c(-1, 1), mean=0, sd=1)
```


```
##
```

```
## If  $X \sim N(0, 1)$ , then
```

```
##  $P(X \leq -1) = P(Z \leq -1) = 0.1587$       $P(X \leq 1) = P(Z \leq 1) = 0.8413$ 
```

```
##  $P(X > -1) = P(Z > -1) = 0.8413$       $P(X > 1) = P(Z > 1) = 0.1587$ 
```

```
##
```




```
## [1] 0.159 0.841
```

and the Think/Show/Tell/Think on pages 402 and 403:

```
sdval <- sqrt(4.50); sdval
```

```
## [1] 2.12
```

```
plotDist("norm", params=list(18, sdval), xlab="x", ylab="f(x)")
```



```
xpnorm(20, mean=18, sd=sdval) # note how exact value is different from the table!
```

```
##
```


```
## If  $X \sim N(18, 2.121)$ , then
```

```
##  $P(X \leq 20) = P(Z \leq 0.9428) = 0.8271$ 
```

```
##  $P(X > 20) = P(Z > 0.9428) = 0.1729$ 
```

```
##
```

■ $P(X \leq 20) = 0.827$ ■ $P(X > 20) = 0.173$



```
## [1] 0.827
```

```
zval <- (20-18)/sdval; zval
```

```
## [1] 0.943
```

```
xpnorm(zval, mean=0, sd=1)
```

```
##
```


```
## If  $X \sim N(0, 1)$ , then
```

```
##  $P(X \leq 0.9428) = P(Z \leq 0.9428) = 0.8271$ 
```

```
##  $P(X > 0.9428) = P(Z > 0.9428) = 0.1729$ 
```

```
##
```

■ $P(X \leq 0.94) = 0.827$ ■ $P(X > 0.94) = 0.173$



```
## [1] 0.827
```